arXiv:1304.6503v1 [math.GT] 24 Apr 2013

LINKS AND SUBMERSIONS TO THE PLANE ON AN OPEN 3-MANIFOLD

SHIGEAKI MIYOSHI

Abstract. We study the realization problem which asks if a given oriented link in an open 3-manifold can be realized as a fibre of a submersion to the Euclidean plane. We correct the results obtained by the author before which contains an error and certain imperfection, and investigate a necessary and sufficient condition for the realization in the words of well-known invariants. We obtain the condition expressed by the first homology group (with mod 2 coefficient).

1. Introduction

The purpose of this note is two-fold. First, we will correct a theorem in [M] by the author, and, second, study a problem which arises from the correction. G. Hector and D. Peralta-Salas found out the error in the theorem in [M] and informed the author about it. Moreover, they studied comprehensively the realization problem which asks if a manifold can be embedded in another manifold so that it is also a fibre of a submersion to the Euclidean space. See [HP]. The correct theorem is the following. Unless otherwise stated, we work in the smooth (C^∞) category in this paper.

Theorem A. *For an oriented link L in an open oriented 3-manifold M, the following conditions are equivalent:*

(1) *there exists a submersion $\varphi : M \to \mathbb{R}^2$ such that up to isotopy the preimage $\varphi^{-1}(0)$ of the origin is L and φ maps the transverse orientation of L to the standard orientation of $\mathbb{R}^2$, i.e., for a small disk D transverse to L with the orientation induced from those of M and L, the restriction $\varphi|D$ preserves the orientation, and*

2000 *Mathematics Subject Classification.* primary 57R30, secondary 57R99, 57M99, 57R25.

Key words and phrases. knot, submersion, completely integrable vector field.

The author is supported by Grant-in-Aid for Scientific Research 23540106.

(2) *the cycle L represents the null-class in the homology of locally finite infinite chains $H_1^\infty(M;\mathbb{Z})$ and a preferred framing of L extends to M.*

For the term "preferred framing", see below. In the original incorrect theorem (Theorem 1 in [M]) the above extension condition of (2) is missing. If (1) holds, then the canonical trivialization of the tangent bundle of $\mathbb{R}^2$ is pulled back and with the tangent vectors of the fibres it determines a framing of the tangent bundle TM which restricts to a preferred framing of L. Conversely, an extension of a preferred framing of L ensures that the projection map of the tubular neighbourhood onto the meridian disk extends to the desired submersion. This is an application of the *h-principle*, i.e., A. Phillips' submersion classification theory [P] in this case. Now we explain this extension condition more precisely and how it fills the gap in the original proof.

First, for an oriented n-component link $L = L_1 \cup L_2 \cup \cdots \cup L_n$ in an open oriented 3-manifold M, we choose a tubular neighbourhood $N(L) \subset M$ and its *framing* $\nu : \coprod_{j=1}^n (S^1 \times D^2)_j \to N(L)$, i.e., an embedding of n solid tori $\coprod_{j=1}^n (S^1 \times D^2)_j$ onto $N(L)$ which takes the cores $\coprod_{j=1}^n (S^1 \times \{0\})_j$ to L. Here, D^2 denotes the unit disk in $\mathbb{C}$ and $S^1 = \partial D^2$. The framing ν and its restriction to the cores should be chosen to be orientation-preserving. Thus, a framing of $N(L)$ is a trivialization of the normal bundle to L realized as a tubular neighbourhood and it also induces a trivialization of the tangent bundle to $N(L)$, $TN(L) = TM|N(L)$. Moreover, in the case that L represents the null-class in $H_1^\infty(M;\mathbb{Z})$, its framing ν of a tubular neighbourhood is said to be *preferred* if the union of the longitudes $\nu\ (\coprod_{j=1}^n (S^1 \times \{1\})_j$ represents the null class in $H_1^\infty(M - \operatorname{Int} N(L);\mathbb{Z})$. Note that a preferred framing is not unique in general even in the case of a knot, unlike in the case of a knot in $\mathbb{R}^3$. We call a framing of a tubular neighbourhood $N(L)$ also a framing of L. The above extension condition of (2) is the requirement of an extension to the whole manifold M of the trivialization of $TM|N(L)$ induced by a preferred framing of $N(L)$. Note that any oriented 3-manifold is parallelizable, i.e., its tangent bundle is trivial.

In the proof in [M], it is only shown that a projection map of the tubular neighbourhood onto a meridian disk with respect to a preferred framing extends to the whole manifold. However, the projection depends on the framing of the tubular neighbourhood and in order to apply Phillips' submersion classification theorem [P] the trivialization of the tubular neighbourhood should extend to the whole manifold. That is, the requirement of the simultaneous extension of the projection map and the framing of the tubular neighbourhood is the necessary and sufficient condition for the existence of the submersion with the properties in (1). See below for the more precise treatment of the extendability of the framing. Theorem A is also a consequence of Theorem 2.4.2 in [HP].

We note that Theorem 2 in [M] is correct even though the proof in [M] is not completed. We state it here for the reader's convenience:

Theorem B ((Theorem 2 in [M]))**.** *For any link L in an open orientable 3-manifold, there is a submersion $\varphi : M \to \mathbb{R}^2$ such that up to isotopy the union of compact components of $\varphi^{-1}(0)$ is L.*

In order to prove this theorem, we have to choose a trivialization of a tubular neighbourhood $N(L)$ which extends to the whole manifold M. This can be always done by twisting a trivialization once around the meridian direction if necessary. Note that we need not to require that the framing is preferred here. This observation is missing in the proof in [M]. Theorem B is also proved in Application 2.3.8 in [HP].

As a consequence of the correction, there arises a question to find a criterion for a link to be a fibre of a submersion in the words of well-known invariants. We will answer to this question for the case of a knot. Suppose that M is an open oriented 3-manifold and K is an oriented knot in M. For the simplicity, we say that K is *realizable* if there is a submersion $\varphi : M \to \mathbb{R}^2$ such that up to isotopy $K = \varphi^{-1}(0)$ and for a small enough disk D transverse to K the restriction $\varphi\,|D$ is orientation-preserving. Here, D is endowed with the orientation induced from those of M and K. Now we fix a trivialization $\Pi : TM \cong M \times \mathbb{R}^3$. Assume that K represents a null-class in the first homology group of locally finite infinite chains. Choose a tubular neighbourhood $N(K)$ of K and a framing $\nu : S^1 \times D^2 \to N(K)$ of $N(K)$. As noted before, the framing ν is an embedding of $S^1 \times D^2$ onto $N(K)$ which is a trivialization of the normal disk bundle to K. Thus, each disk $\nu\left(\{\exp(t\sqrt{-1})\} \times D^2\right)$ may be assumed to be normal to K with respect to the metric induced by the parallelization Π. Then, the orientations of M and K, the parallelization Π, and the framing ν altogether induce a map $f : K \to \mathrm{SO}(3)$. More precisely, for any $p \in K$ we set $v_1(p) = \dfrac{D\nu\frac{\partial}{\partial\theta}}{\|D\nu\frac{\partial}{\partial\theta}\|}$ and $v_2(p) = \dfrac{D\nu\frac{\partial}{\partial x}}{\|D\nu\frac{\partial}{\partial x}\|}$, where we write $z = x + y\sqrt{-1} \in D^2$, and choose $v_3(p) \in T_pM$ so that $(v_1(p), v_2(p), v_3(p))$ defines an orthonormal frame of T_pM. With respect to Π, this orthonormal frame $(v_1(p), v_2(p), v_3(p))$ can be expressed as a special orthogonal matrix which is defined to be $f(p) \in \mathrm{SO}(3)$. We call this map $f : K \to \mathrm{SO}(3)$ an *h-framing* of K. Note that $[S^1, \mathrm{SO}(3)] \cong \mathrm{Hom}(\pi_1(S^1), \pi_1(\mathrm{SO}(3))) \cong \pi_1(\mathrm{SO}(3)) \cong \mathbb{Z}_2$ with appropriate choices of base points. Thus, we may identify the homotopy class $[f]$ of an h-framing with its image $f_*[K] \in \pi_1(\mathrm{SO}(3))$. A framing of a knot K extends to the ambient manifold M if and only if its h-framing extends to M.

Next, we set $\mu = \nu(\{1\} \times \partial D^2)$, the meridian loop on $\partial N(K)$. Let $\langle\mu\rangle^\infty$ denote the subgroup of $H_1^\infty(E_M(K);\mathbb{Z})$ generated by $[\mu]$, where $E_M(K)$ denotes the exterior of K in M, i.e., $E_M(K) := M - \mathrm{Int}\, N(K)$. Note that $\langle\mu\rangle^\infty$ is isomorphic to $\mathbb{Z}_n = \mathbb{Z}/n\mathbb{Z}$ for some $n \in \mathbb{Z}_{\geq 0}$, where we consider $\mathbb{Z}_0$ as $\mathbb{Z}$ and $\mathbb{Z}_1$ as 0. Now we state an answer to the above question.

Theorem 1. *Assume that K represents the null-class in $H_1^\infty(M;\mathbb{Z})$. If $\langle\mu\rangle^\infty$ is isomorphic to $\mathbb{Z}_{2m+1}$ ($m \in \mathbb{Z}_{\geq 0}$), then K is realizable.*

Theorem 2. *Assume that K represents the null-class in $H_1^\infty(M;\mathbb{Z})$ and $H_1(M;\mathbb{Z})$ is finitely generated. If $\langle\mu\rangle^\infty$ is isomorphic to $\mathbb{Z}_{2m}$ ($m \in \mathbb{Z}_{\geq 0}$), then K is realizable if and only if K represents non-zero class in $H_1(M;\mathbb{Z}_2)$.*

Remark 1. It can be seen by a simple construction that there is a knot K in an open orientable 3-manifold M such that one of the conditions of Theorem 1 and 2 is satisfied respectively. As one of the simplest examples, suppose that $M = S^1 \times \mathrm{Int}\, D^2$ and K is the $(p, 1)$ cable knot of the core circle of M (see

Section 3 for the precise description of cable knots). Obviously K represents the null-class in $H_1^\infty(M;\mathbb{Z}) \cong H^2(M;\mathbb{Z}) = 0$ and $\langle\mu\rangle^\infty \subset H_1^\infty(E_M(K);\mathbb{Z})$ is isomorphic to $\mathbb{Z}_p$. In the case that p is odd, it meets Theorem 1. In this case, one can construct a submersion $M \to \mathbb{R}^2$ concretely which shows the realizability of K. The construction may show to some extent that the reason why the parity of p rules the realizability. In the case that p is even, K and M meet the half of Theorem 2, i.e., $[K] = 0$ in $H_1(M;\mathbb{Z}_2)$ and $\langle\mu\rangle^\infty \cong \mathbb{Z}_p$ (p : even). An example which meets the other half of Theorem 2 can be constructed with the handlebody of genus $2m$ by performing a Dehn surgery, for example. Anyway, there is a knot K in an open orientable 3-manifold M such that $[K] = 0$ in $H_1^\infty(M;\mathbb{Z})$, $\langle\mu\rangle^\infty \cong \mathbb{Z}_{2m}$ and $[K] \neq 0$ in $H_1(M;\mathbb{Z}_2)$. A concrete construction not by using h-principle but by hand must be interesting itself and deepen our understanding.

Remark 2. If the meridian of K is odd-torsion in $H_1^\infty(M;\mathbb{Z})$ then the homology class $[K] \in H_1(M;\mathbb{Z})$ is not an even-multiple of any class. This fact can be easily seen by considering the intersection form $H_1(M;\mathbb{Z}) \otimes H_2^\infty(M;\mathbb{Z}) \to \mathbb{Z}$.

Remark 3. As mentioned earlier, G. Hector and D. Peralta-Salas [HP] studied this kind of realization problem in the more general dimensions and setting. As one application of their theory, they obtained a characterization for a link in $\mathbb{R}^3$ to be realizable and in particular they showed that no knot in $\mathbb{R}^3$ is realizable. One may consider Theorem 2 generalizes the result.

Remark 4. In the case of links, the argument will be a rather complicated nuisance. It might be just a technicality, nevertheless we omit here the consideration in the case of links at all. The complete research including the general case of links should be done in the sequel.

2. Proof of Theorem 1

First, we need the following lemma.

Lemma 1. *Assume that K represents the null-class in $H_1^\infty(M;\mathbb{Z})$. Then the following hold.*

(1) *If* $\langle\mu\rangle^\infty$ *is isomorphic to* $\mathbb{Z}_{2m+1}$ $(m \in \mathbb{Z}_{\geq 0})$*, then there are preferred framings* $\nu_j : S^1 \times D^2 \to N(K)$ $(j = 0, 1)$ *which induce h-framings* $f_j : K \to \mathrm{SO}(3)$ *such that* $[f_j] = j \in \mathbb{Z}_2 \cong \pi_1(\mathrm{SO}(3))$.

(2) *If* $\langle\mu\rangle^\infty$ *is isomorphic to* $\mathbb{Z}_{2m}$ $(m \in \mathbb{Z}_{\geq 0})$*, then the h-framing induced by any preferred framing is unique up to homotopy.*

Proof. (1) Suppose $\nu : S^1 \times D^2 \to N(K)$ is a preferred framing and $f_\nu : K \to \mathrm{SO}(3)$ is the h-framing induced by ν. Let λ denote the longitude $\nu(S^1 \times \{1\})$ and recall $\mu = \nu(\{1\} \times \partial D^2)$. Thus, the kernel of the homomorphism $H_1^\infty(\partial N(K); \mathbb{Z}\ ) \to H_1^\infty(E_M(K); \mathbb{Z}\ )$ induced by the inclusion $\partial N(K) \hookrightarrow E_M(K) = M - \mathrm{Int}\, N(K)$ is generated by $[\lambda]$ and $(2m+1)[\mu]$. Therefore $[\lambda] + n(2m+1)[\mu] = 0 \in H_1^\infty(E_M(K); \mathbb{Z}\ )$ for any $n \in \mathbb{Z}$. Now, let ν' be the framing $S^1 \times D^2 \to N(K)$ obtained by twisting the preferred framing ν $2m+1$ times around μ-direction. Then ν' is also preferred and $[f_{\nu'}] = [f_\nu] + (2m+1) \not\equiv [f_\nu] \in \pi_1(\mathrm{SO}(3))$, where we consider $\pi_1(\mathrm{SO}(3))$ as $\mathbb{Z}_2$. This shows (1).

(2) As in the proof of (1), $[\lambda] + n(2m)[\mu] = 0 \in H_1^\infty(E_M(K); \mathbb{Z}\ )$ for any $n \in \mathbb{Z}$. We fix a preferred framing $\nu : S^1 \times D^2 \to N(K)$. For any other preferred framing $\nu' : S^1 \times D^2 \to N(K)$ and its induced h-framing $f_{\nu'}$, we have $[f_{\nu'}] = [f_\nu] + n(2m) \equiv [f_\nu] \in \pi_1(\mathrm{SO}(3))$. Thus the h-framing induced by a preferred framing is unique (up to homotopy).

□

In other words, we may say that if μ is even-torsion in $H_1^\infty(E_M(K); \mathbb{Z}\ )$ then the *preferred h-framing*, i.e., the h-framing induced by a preferred framing is unique and if μ is odd-torsion then we can realize any preferred h-framing by choosing an appropriate preferred framing. Now, we proceed to the proof of Theorem 1.

Proof of Theorem 1. Assume that $\langle\mu\rangle^\infty$ is isomorphic to $\mathbb{Z}_{2m+1}$ $(m \in \mathbb{Z}_{\geq 0})$. Then, by (1) of Lemma 1 we have a preferred framing $\nu : S^1 \times D^2 \to N(K)$ which induces an h-framing $f_\nu : K \to \mathrm{SO}(3)$ with $[f_\nu] = 0$ in $\pi_1(\mathrm{SO}(3))$. Thus, trivial homomorphism $\pi_1(M) \to \pi_1(\mathrm{SO}(3))$ is an extension of $(f_\nu)_* :$

$\pi_1(K) \to \pi_1(\mathrm{SO}(3))$. Note that M is homotopy equivalent to a subcomplex of its 2-skeleton with respect to a triangulation of M (cf. [W], for example). Therefore, the existence of an extension of the homomorphism $(f_\nu)_* : \pi_1(K) \to \pi_1(\mathrm{SO}(3))$ to $\pi_1(M)$ implies the existence of an extension of the map $f_\nu : K \to \mathrm{SO}(3)$ to M. Consequently, by Theorem A, K is realizable. □

As we see in the proof of Theorem 1, we have a criterion on the existence of an extension of a framing of K to M. As mentioned in the proof of Theorem 1, an open orientable 3-manifold is homotopy equivalent to a subcomplex of its 2-skeleton. Therefore, by an elementary obstruction theory, a map $f : K \to \mathrm{SO}(3)$ extends to a map $M \to \mathrm{SO}(3)$ if and only if the induced homomorphism $f_* : \pi_1(K) \to \pi_1(\mathrm{SO}(3))$ extends to a homomorphism $\pi_1(M) \to \pi_1(\mathrm{SO}(3))$, i.e., there is a homomorphism $\Phi : \pi_1(M) \to \pi_1(\mathrm{SO}(3))$ such that $\Phi \circ \iota_* = f_*$ where $\iota : K \hookrightarrow M$ is the inclusion. Moreover, since $\pi_1(K)$ and $\pi_1(\mathrm{SO}(3))$ are Abelian it is equivalent to the condition that the homomorphism $f_* : H_1(K;\mathbb{Z}) \to H_1(\mathrm{SO}(3);\mathbb{Z})$ extends to a homomorphism $H_1(M;\mathbb{Z}) \to H_1(\mathrm{SO}(3);\mathbb{Z})$. Thus, we have the following.

Lemma 2. *Let M be an open orientable 3-manifold and K a knot in M. Then the following are equivalent:*

1. *a map $f : K \to \mathrm{SO}(3)$ extends to a map $M \to \mathrm{SO}(3)$,*
2. *the induced homomorphism $f_* : \pi_1(K) \to \pi_1(\mathrm{SO}(3))$ extends to a homomorphism $\pi_1(M) \to \pi_1(\mathrm{SO}(3))$, and*
3. *the homomorphism $f_* : H_1(K;\mathbb{Z}) \to H_1(\mathrm{SO}(3);\mathbb{Z})$ extends to a homomorphism $H_1(M;\mathbb{Z}) \to H_1(\mathrm{SO}(3);\mathbb{Z})$.*

3. A key lemma to the proof of Theorem 2

In this section, we will study the case of (2) of Lemma 1 and show a key lemma to the proof of Theorem 2. In this case, the preferred h-framing of the knot K is unique (up to homotopy), which makes the situation restrictive and more complicated. In order to state the key lemma, we define the notion of cabling a knot in a manifold and its natural framing. Suppose that p is a positive integer and q is an integer. Set $\theta = \frac{2\pi}{p}$. We define p parallel line $\tilde{L}$ in $\mathbb{R} \times D^2$ by

$$\tilde{L} := \{(t, \tfrac{1}{2}\exp(k\theta\sqrt{-1})) \mid t \in \mathbb{R}, k \in \mathbb{Z}, 0 \leq k < p\}$$

and a self-diffeomorphism $\rho_{p/q}$ on $\mathbb{R}\times D^2$ by $\rho_{p/q}(t,z) := (t+1, \exp(-q\theta\sqrt{-1})z)$. The diffeomorphism $\rho_{p/q}$ induces an action on $\mathbb{R} \times D^2$ and its quotient $(\mathbb{R}\times D^2)/\rho_{p/q}$ is diffeomorphic to $S^1 \times D^2$. Precisely, let $\tilde{\eta} : \mathbb{R}\times D^2 \to \mathbb{R}\times D^2$ be a diffeomorphism defined by $\tilde{\eta}(t,z) := (t, \exp(-q\theta t\sqrt{-1})z)$. Then $\tilde{\eta}$ is an equivariant with respect to the translation and $\rho_{p/q}$, i.e., $\rho_{p/q} \circ \tilde{\eta}(t,z) = \tilde{\eta}(t+1, z)$ for any $(t,z) \in \mathbb{R} \times D^2$. Hence it induces a diffeomorphism $\eta : (\mathbb{R}/\mathbb{Z}) \times D^2 \to (\mathbb{R} \times D^2)/\rho_{p/q}$. Since $\exp(2\pi t\sqrt{-1}) \mapsto t \bmod \mathbb{Z}$ is a diffeomorphism from S^1 to $\mathbb{R}/\mathbb{Z}$, we have a desired diffeomorphism from $S^1 \times D^2$ to $(\mathbb{R} \times D^2)/\rho_{p/q}$. Let $L_{p,q}$ denote the quotient of $\tilde{L}$ by this action. Then $L_{p,q}$ is a link in $S^1 \times D^2$ and called a (p,q) *cable link* of the core curve $S^1 \times \{0\}$. In fact, for the standard solid torus $S^1 \times D^2$ in S^3, $L_{p,q}$ is just the (p,q) torus link in S^3. For any knot K_0 in a manifold, we choose a framing $\nu\ : S^1 \times D^2 \to N(K_0)$ of a tubular neighbourhood and the image $\nu\ (L_{p,q})$ is defined to be a (p,q) cable link of K_0. Note that it depends on the choice of the framing of $N(K_0)$.

To define a natural framing of a cable link, we consider annuli in $S^1 \times D^2$ defined as follows. Let $\tilde{A}$ be p strips in $\mathbb{R} \times D^2$ defined by

$$\tilde{A} := \{(t, r\exp(k\theta\sqrt{-1})) \mid t \in \mathbb{R}, \tfrac{1}{2} \leq r \leq 1, k \in \mathbb{Z}, 0 \leq k < p\}$$

and set $A_{p,q} := \tilde{A}/\rho_{p/q}$, the quotient of $\tilde{A}$ by the action of the diffeomorphism $\rho_{p/q}$. Thus $A_{p,q}$ is a union of annuli in $S^1 \times D^2$ bounded by $L_{p,q}$ and a union of mutually disjoint parallel curves in $S^1 \times \partial D^2$. For any small tubular neighbourhood $N(L_{p,q})$, the section $A_{p,q} \cap N(L_{p,q})$ determines a framing of $N(L_{p,q})$. We call this framing the *revolution framing* of the cable link $L_{p,q}$. See Figure 1. Note that if p and q are coprime then $L_{p,q}$ is connected and called a (p,q) cable *knot*.

The following is the key lemma. By this lemma, we transport the given situation on K to a certain neighbourhood of a basic knot for K.



Figure 1. A revolution framing of $L_{p,q}$ at a section

Lemma 3. *Let J be an oriented knot in M which represents the null-class in $H_1^{\infty}(M;\mathbb{Z})$. Suppose $[J] = p\gamma$ where p is a positive integer and $\gamma \in H_1(M;\mathbb{Z})$. Fix a representative cycle C for γ: $[C] = \gamma$. Then there exists a cable knot L of C of type $(p,1)$ if p is even and of type $(p,1)$ or $(p,2)$ if p is odd which has the following properties:*

1. *L is homologous to J,*
2. *a preferred framing of J induces a preferred framing of L, and*
3. *the h-framing of L determined by the preferred framing of L is equal to a revolution h-framing of C.*

Proof. Since $[J] = p[C]$ in $H_1(M;\mathbb{Z})$, there is a chain bounded by J and p times of C. Precisely, there is an immersion from a compact orientable surface S into M with the following properties. The boundary ∂S decomposes into a component $\partial_+ S$ and a union $\partial_- S$ of components. The immersion is an embedding except $\partial_- S$ where the immersion is a p-fold covering map onto C. Moreover, $\partial_+ S$ is mapped diffeomorphically onto J. Thus the image is a compact (singular) surface in M which is non-singular away from C and in a small neighbourhood of C it is a mapping cylinder of a finite covering map $\partial_- S \to C$. By abus of notation, let S denote the image of the immersion $S \to M$. We choose a small enough tubular neighbourhood $N(C)$. In general position, $\partial N(C)$ and S are transverse to each other and we may assume that $S \cap \partial N(C)$ is a union of parallel, cooriented circles by performing double curve surgery on S (and $\partial N(C)$) if necessary. Thus we may assume that the covering map $\partial_- S \to C$ is p-fold. Moreover, by choosing appropriate framing of $N(C)$, we may also assume that $S \cap \partial N(C)$ is a (p, q) cable link of C, where $0 \leq q < p$.

We will deform S in $N(C)$ by performing double curve surgery as follows. Consider the concentric tubular neighbourhood $N_{1/2}(C) \subset N(C)$ where normal disks are of radius $\frac{1}{2}$ of the normal disks of $N(C)$. We will set a cable knot L on $\partial N_{1/2}(C)$ and construct a compact oriented surface B in $N(C)$ bounded by $S \cap \partial N(C)$ and L.



Figure 2. Double curve surgery on A and meridian disks

Now suppose that p is even. We set L to be the $(p, 1)$ cable knot on $\partial N_{1/2}(C)$ and choose a small tubular neighbourhood $N(L)$ of L. Set $A = A_{p,1} - \operatorname{Int} N(L)$, where $A_{p,1}$ is the annulus in $N(C)$ bounded by $L = L_{p,1}$ and a curve on $\partial N(C)$ defined earlier. The orientation of A is naturally determined so that it induces the given orientation of L. Moreover, consider disjoint $q-1$ copies of the meridian disk in $N(C)$ with the natural orientation induced by those of C and M. Here, in the case $q = 0$, -1 copies of the meridian disk means one meridian disk with the opposite orientation. Performing double curve surgery on the union of A and these meridian disks yields a compact proper surface B in $N(C) - \operatorname{Int} N(L)$ so that $B \cap \partial N(C)$ is a (p, q) cable link and $B \cap \partial N(L)$ is isotopic to L in $N(L)$. See Figure 2 and 3. Note that the number of the intersection points of L and $q-1$ meridian disks is $p|q-1|$, an even number.



(a) (6, 1) cable and two meridian loops

(b) (6, 3) cable

FIGURE 3. The effect on a section: $p = 6$ and $q = 3$

Next, suppose that p is odd. If q is even, then we set L to be the $(p, 2)$ cable knot on $\partial N_{1/2}(C)$ and A to be the oriented spanning annulus $A_{p,2} - \operatorname{Int} N(L)$ similarly to the case p is even. Consider $q-2$ copies of the oriented meridian disk and perform double curve surgery on A and these meridian disks to obtain the desired surface B. As above, if n is an negative integer, then n copies means $|n|$ copies with the opposite orientation. If q is odd, then we set L to be the $(p, 1)$ cable knot on $\partial N_{1/2}(C)$ and A to be the spanning annulus $A_{p,1} - \operatorname{Int} N(L)$. This time we perform double curve surgery on A and $q-1$ meridian disks to obtain the desired surface B. Note that the number of the intersection points of L and the meridian disks are $p|q-2|$ if q is even and $p|q-1|$ if q is odd, and both of them are even.

In any case above, we have a compact oriented proper surface B in $N(C) - \operatorname{Int} N(L)$ such that $B \cap \partial N(L)$ is isotopic to L in $N(L)$ and $B \cap \partial N(C)$ is isotopic to $S \cap \partial N(C)$ in $\partial N(C)$. Therefore we may consider that B is bounded by $S \cap \partial N(C)$ and L. Thus the surface $S - \operatorname{Int} N(C)$ attached to B along $S \cap \partial N(C) = B \cap \partial N(C)$ is bounded by J and L. We define a framing of L by setting the curve $B \cap N(L)$ to be the longitude.

As for the h-framing of L, the effect of the surgery is to twist once at each intersection point of L with meridian disks. See Figure 2. As noted above, the number of the intersection is even in any case, which implies the h-framing of L determined by the surface B (i.e., induced from the preferred h-framing $f : J \to \mathrm{SO}(3)$) is equal to the h-framing defined by the revolution framing. This completes the proof of Lemma 3. □

4. Proof of Theorem 2

In this last section, we will prove Theorem 2. In view of Theorem A and Lemma 2, only the extendability of the homomorphism on the first homology groups will be our concern. Lemma 3 will be used in the both direction in the proof. To show the condition is necessary (Assertion 1 in the proof below), we will describe the homological situation with a preferred framing by Lemma 3. To show the condition is sufficient (Assertion 2 in the proof below), we will use Lemma 3 to adjust a preferred framing and define an extended homomorphism.

Proof of Theorem 2. First, we fix a direct sum decomposition of $H_1(M;\mathbb{Z})$:

$$H_1(M;\mathbb{Z}) = \mathrm{Fr}(H_1(M;\mathbb{Z})) \oplus \mathrm{Tor}(H_1(M;\mathbb{Z}))$$

where $T := \mathrm{Tor}(H_1(M;\mathbb{Z}))$ denotes the torsion subgroup and $F := \mathrm{Fr}(H_1(M;\mathbb{Z})) \cong H_1(M;\mathbb{Z})/T$. For the simplicity, we write $\kappa = [K] \in H_1(M;\mathbb{Z})$ and with respect to the above direct sum decomposition, we write $\kappa = \kappa_F + \kappa_T \in F \oplus T$. By the assumption of the theorem and Lemma 1, a preferred h-framing of K is unique up to homotopy.

By Theorem A and Lemma 2, we only have to show the following two assertions. Let $\zeta_{(2)}$ denote the $\mathbb{Z}_2$-reduction in $H_1(M;\mathbb{Z}_2)$ of a class $\zeta \in H_1(M;\mathbb{Z})$.

Assertion 1. If $\kappa_{(2)} = 0$, then f_* does not extend to a homomorphism $H_1(M;\mathbb{Z}) \to H_1(\mathrm{SO}(3);\mathbb{Z})$.

Assertion 2. If $\kappa_{(2)} \neq 0$, then f_* extends to a homomorphism $H_1(M;\mathbb{Z}) \to H_1(\mathrm{SO}(3);\mathbb{Z})$.

We note that $\kappa_{(2)} = 0$ if and only if $(\kappa_F)_{(2)} = 0 \in H_1(M;\mathbb{Z}_2)$ and either κ_T is odd-torsion or an even multiple of an even-torsion class in $H_1(M;\mathbb{Z})$.

Proof of Assertion 1. Suppose on the contrary to the conclusion there is an extension of $f_* : H_1(K;\mathbb{Z}) \to H_1(\mathrm{SO}(3);\mathbb{Z})$ to a homomorphism $\Phi : H_1(M;\mathbb{Z}) \to H_1(\mathrm{SO}(3);\mathbb{Z})$. Then since $\kappa_{(2)} = 0$ and $H_1(\mathrm{SO}(3);\mathbb{Z}) \cong \mathbb{Z}_2$, we have $f_*([K]) = \Phi \circ \iota_*([K]) = \Phi(\kappa) = 0$.

Since $(\kappa_F)_{(2)} = 0$, there are a primitive class $\alpha \in H_1(M;\mathbb{Z})$ and a non-negative integer m such that $\kappa_F = 2m\alpha$. Here, a homology class ζ is *primitive* if there is no class whose any multiple is equal to ζ. As noted above, $(\kappa_T)_{(2)} = 0$ if and only if κ_T is odd-torsion or an even-multiple of an even torsion class. Thus, we have two cases.

Case 1. Assume that κ_T is odd-torsion. Suppose that the order of κ_T is k. Then $\kappa_T = (k-1)(-\kappa_T) = 2n(-\kappa_T)$ for some non-negative integer n. Hence, $\kappa = \kappa_F + \kappa_T = 2m\alpha + 2n(-\kappa_T) = 2(m\alpha + n(-\kappa_T))$. Setting $J = K, \gamma = m\alpha + n(-\kappa_T)$ and $p = 2$, we apply Lemma 3. Then we have the $(2,1)$ cable knot L of a representative cycle C of γ with respect to a framing of $N(C)$. Denote the induced h-framing by $g : C \to \mathrm{SO}(3)$ and the revolution h-framing $f_L : L \to \mathrm{SO}(3)$. In $H_1(\mathrm{SO}(3);\mathbb{Z}) \cong \mathbb{Z}_2$ we have $[f_L] = 2[g]+1 \equiv 1$, which implies the h-framing of L is non-zero whatever the framing of C may be. Since $[L] = \kappa$, we have $0 = \Phi(\kappa) = \Phi([L]) = [f_L] \equiv 1$ in $H_1(\mathrm{SO}(3)\mathbb{Z})$. This is a contradiction.

Case 2. Assume that κ_T is an even multiple of an even-torsion class. Since $\kappa_T = 2n\beta$ for some $\beta \in H_1(M;\mathbb{Z})$ and a non-negative integer n, we have $\kappa = \kappa_F + \kappa_T = 2(m\alpha + n\beta)$. Applying Lemma 3 to $J = K, \gamma = m\alpha + n\beta$ and $p = 2$, we have a contradiction similarly as in Case 1.

Thus we have a contradiction in any case, which implies that f_* never extends to $H_1(M;\mathbb{Z})$. This proves Assertion 1. ☐

Proof of Assertion 2. Assume that $\kappa_{(2)} \neq 0$, i.e., $(\kappa_F)_{(2)} \neq 0$ or $(\kappa_T)_{(2)} \neq 0$ in $H_1(M;\mathbb{Z}_2)$. The latter is equivalent to the condition that κ_T is an odd multiple of a non-zero even-torsion class in $H_1(M;\mathbb{Z})$. We will construct a homomorphism $\Phi : H_1(M;\mathbb{Z}) \to H_1(\mathrm{SO}(3);\mathbb{Z})$ which is an extension of

$f_* : H_1(K;\mathbb{Z}) \to H_1(\mathrm{SO}(3);\mathbb{Z})$. We choose representative cycles K_F and K_T for the classes κ_F and κ_T respectively. Moreover, a preferred framing of K induces natural framings of K_F and K_T which induce h-framings $f_{K_F} : K_F \to \mathrm{SO}(3)$ and $f_{K_T} : K_T \to \mathrm{SO}(3)$ respectively.

To construct the desired homomorphism, we will work on summand by summand. First, we assume that $(\kappa_F)_{(2)} \neq 0$ and $(\kappa_T)_{(2)} \neq 0$.

Step 1. Since $(\kappa_F)_{(2)} \neq 0$, there are a (torsion-free) primitive class $\alpha \in H_1(M;\mathbb{Z})$ and a non-negative integer m such that $\kappa_F = (2m+1)\alpha$. Apply Lemma 3 to $J = K_F$, $\gamma = \alpha$ and $p = 2m+1$. Then we have a $(2m+1, 1)$ or $(2m+1, 2)$ cable knot L of a representative C of α with respect to a framing of $N(C)$. Suppose $g : C \to \mathrm{SO}(3)$ is the induced h-framing. Since $[L] = [K_F]$, in order to obtain an extension we have to adjust the revolution h-framing f_L so that $[f_L] = [f_{K_F}]$. Now in $H_1(\mathrm{SO}(3);\mathbb{Z})$ we have $[f_L] = (2m+1)[g] + r$, where $r = 1$ or 2 depending on the type of the cable knot L. If $r = 1$, then we twist g once in the direction of meridian of $N(C)$ to have another h-framing $g' : C \to \mathrm{SO}(3)$. As for g', we have $[f_L] = (2m+1)([g'] - 1) + 1 \equiv [g'] \in \mathbb{Z}_2 \cong H_1(\mathrm{SO}(3);\mathbb{Z})$. In this case, set $h = g'$. If $r = 2$, then $[f_L] = (2m+1)[g] + 2 \equiv [g]$ and we set $h = g$. Since by the assumption of the theorem, we can choose a basis $\{\alpha_1, \alpha_2, \ldots, \alpha_b\}$ of F extending $\alpha = \alpha_1$. Define $\Phi_F : H_1(M;\mathbb{Z}) \to H_1(\mathrm{SO}(3);\mathbb{Z})$ as follows:

$$\begin{aligned} \Phi_F(\alpha_i) &= \begin{cases} [h] & \text{if } i = 1 \\ 0 & \text{otherwise} \end{cases} \\ \Phi_F|T &\equiv 0 \end{aligned}$$

Then Φ_F is a well-defined extension of $(f_{K_F})_*$.

Step 2. Recall that $(\kappa_T)_{(2)} \neq 0$ if and only if κ_T is an odd-multiple of a non-zero even-torsion class. We define a subgroup G of T as follows. Let $\langle \kappa_T \rangle$ denote the subgroup generated by κ_T in T. We define a subset $D^{\mathrm{odd}}(\kappa_T)$ of T by

$$D^{\mathrm{odd}}(\kappa_T) := \{\sigma \in T \mid \kappa_T = (2m+1)\sigma \text{ for some } m \in \mathbb{Z}\},$$

and we define G to be the subgroup generated by $D^{\mathrm{odd}}(\kappa_T)$. Note that $\langle \kappa_T \rangle$ is a subgroup of G. Then by the fundamental theorem for finitely generated

Abelian groups, it is easily seen that G is a direct summand of T (hence, of $H_1(M;\mathbb{Z})$). We define a homomorphism $\Phi_T : G \to H_1(\mathrm{SO}(3);\mathbb{Z})$ by setting $\Phi_T|D^{\mathrm{odd}}(\kappa_T) \equiv (f_{K_T})_*[K_T]$. It can be seen that this is well-defined. We extend Φ_T to $H_1(M;\mathbb{Z})$ as zero homomorphism on the complementary summand of G in $H_1(M;\mathbb{Z})$, which is an extension of $(f_{K_T})_*$.

Now we have extensions Φ_F and Φ_T of $(f_{K_F})_*$ and $(f_{K_T})_*$ respectively. Hence setting $\Phi = \Phi_F + \Phi_T$, we have

$$\begin{aligned} \Phi \circ \iota_*[K] &= (\Phi_F + \Phi_T) \circ \iota_*[K] \\ &= \Phi_F \circ \iota_*[K_F] + \Phi_T \circ \iota_*[K_T] \\ &= (f_{K_F})_*[K_F] + (f_{K_T})_*[K_T] \\ &= f_*[K]. \end{aligned}$$

Thus, we have the desired extension Φ of f_*.

In the case that $(\kappa_F)_{(2)} = 0$ or $(\kappa_T)_{(2)} = 0$, we have to modify the above construction in order to adjust the homology class. Indeed, if $(\kappa_F)_{(2)} = 0$, then $(\kappa_T)_{(2)} \neq 0$ and we define Φ_T by setting $\Phi_T|D^{\mathrm{odd}}(\kappa_T) \equiv f_*[K]$ and $\Phi_F \equiv 0$. If $(\kappa_T)_{(2)} = 0$, then $(\kappa_F)_{(2)} \neq 0$ and we define $\Phi_F(\alpha_1) = f_*[K](= [h] + (f_{K_T})_*[K_T])$ and $\Phi_T \equiv 0$. Then $\Phi = \Phi_F + \Phi_T$ is the desired extended homomorphism in these cases. Thus we obtain Assertion 2. □

Consequently, the proof of Theorem 2 is completed. □

Remark 5. The assumption that $H_1(M;\mathbb{Z})$ is finitely generated is used in constructing an extended homomorphism, in particular in Step 2 in the proof of Assertion 2. As the proof shows, the condition $\kappa_{(2)} \neq 0$ is necessary even in the infinitely generated case. The question which asks if $\kappa_{(2)} \neq 0$ is sufficient in general is still open.

Acknowledgements. The author would like to express his hearty gratitude to Gilbert Hector who informed him that his earlier work contains an error. He also thanks Daniel Peralta-Salas for studying, with G. Hector, the problem which he concerned before and is left caused by his misunderstanding. Without their notice the author would never obtain the correction and new results in the point of the author's view.

References

[HP] **G. Hector, D. Peralta-Salas**, Integrable embeddings and foliations, *Amer. J. Math.* 134 (2012), 773–825.

[M] **S. Miyoshi**, Links and globally completely integrable vector fields on an open 3-manifold, *Topology* 34 (1995) 383–387.

[P] **A. Phillips**, Submersions of open manifolds, *Topology* 6 (1966) 171–295.

[W] **J. H. C. Whitehead**, The immersion of an open 3-manifold in Euclidean 3-space, *Proc. London Math. Soc.* 11 (1961) 81–90.

DEPARTMENT OF MATHEMATICS, CHUO UNIVERSITY, 1-13-27 KASUGA BUNKYO-KU, TOKYO, 112-8551, JAPAN

E-mail address: `miyoshi@math.chuo-u.ac.jp`